

A DISTRIBUTIVE LATTICE COVER FOR SEMILATTICES

COLIN G. BAILEY AND JOSEPH S. OLIVEIRA

ABSTRACT. We consider two constructions of an envelope for a finite locally distributive strong upper semilattice. The first is based on Birkhoff's representation of finite distributive lattices and the second on valuations on lattices. We show that these produce isomorphic envelopes.

1. INTRODUCTION

There are many interesting semilattices that also have meets whenever lower bounds exist – see [1–3] for some implication based examples. For convenience we refer to these as strong upper semilattices. Thus these structures have the property that every interval $[a, b]$ is a lattice, and if $[a, b] \subseteq [c, d]$ then $[a, b]$ is a sublattice of $[c, d]$.

We consider those strong semilattices in which every interval is distributive. In this paper we concentrate primarily on finite examples and show that two well-known constructions lead to an *envelope*, ie a distributive lattice in which our original structure embeds as an upwards-closed sub-semilattice.

The first construction is based upon the Birkhoff representation theorem for finite distributive lattices, and the second is based upon the valuation ring of a lattice. We show that these two constructions lead to the same (up to isomorphism) envelope.

Definition 1.1. A strong upper semilattice is an upper semilattice with one such that any two elements with a lower bound have a greatest lower bound. As usual we will denote this glb by $x \wedge y$.

Definition 1.2. A distributive strong upper semilattice is a strong upper semilattice in which every interval $[a, b]$ is a distributive lattice.

2. THE BIRKHOFF CONSTRUCTION

Throughout this section we assume that $\mathcal{L}$ is a **finite** distributive strong upper semilattice. We intend to use the meet-irreducibles of $\mathcal{L}$ to get a distributive lattice as is done in Birkhoff's representation theorem for finite distributive lattices. Then we show that our original semilattice embeds into this new lattice as an upwards-closed subalgebra.

The first part of this proof verifies that the usual machinery for representation by meet-irreducibles works in our setting.

Let $\mathcal{M}$ be the set of meet-irreducibles of $\mathcal{L}$. and if $x \in \mathcal{L}$ then $\mathcal{M}^x = \{m \in \mathcal{M} \mid x \leq m\}$.

Let $A = \{a \in \mathcal{L} \mid a \text{ is minimal}\}$.

Lemma 2.1.

$$x = \bigwedge \mathcal{M}^x.$$

Date: February 2, 2009.

1991 Mathematics Subject Classification. 06A12, 06D05.

Key words and phrases. semilattice, distributive lattice, valuation.

Proof. By induction on co-rank. Everything of corank one is in $\mathcal{M}$ and so the result is clear.

Obviously $x \leq \bigwedge \mathcal{M}^x = s$. Suppose that $x < s$. Now if $x = p \wedge q$ for any $x < p, q$ in $\mathcal{L}$ then $p = \bigwedge \mathcal{M}^p$ and $q = \bigwedge \mathcal{M}^q$ and $\mathcal{M}^p \cup \mathcal{M}^q \subseteq \mathcal{M}^x$ so that $x = p \wedge q = \bigwedge (\mathcal{M}^p \cup \mathcal{M}^q) \geq \bigwedge \mathcal{M}^x > x$ – contradiction. Hence x must be in $\mathcal{M}$ and so the result is clear. $\square$

Lemma 2.2. *Let $x \in \mathcal{M}$ and $x^+ = \bigwedge \{y \in \mathcal{M} \mid x < y\}$. Then*

- i. $x < x^+$;
- ii. if $z \in \mathcal{M}$ is not equal to x then $x \vee z = x^+ \vee z$.
- iii. if $z \in \mathcal{M}$ is not equal to x then $x \vee z = x^+ \vee z^+$.

Proof. i. This is immediate as x is meet-irreducible.

ii. If $z \in \mathcal{M}^x$ then $z \geq x^+$ and both sides give z . So we may assume that $z \notin \mathcal{M}^x$.

Obviously $x \vee z \leq x^+ \vee z$. If $y \in \mathcal{M}^{x \vee z}$ then $y > x$ and so $y \geq x^+$. Hence $y \geq x^+ \vee z$.

Thus $x \vee z = \bigwedge \mathcal{M}^{x \vee z} = \bigwedge \mathcal{M}^{x^+ \vee z} = x^+ \vee z$.

iii. $x \neq z$ gives $x \vee z = x^+ \vee z = x \vee z^+$. Hence $x^+ \vee z^+ = (x \vee x^+) \vee (z \vee z^+) = (x \vee z^+) \vee (x^+ \vee z) = x \vee z$. $\square$

2.1. Using Filters. Now for the heart of the proof – we let $\mathcal{D} = \mathcal{D}(\mathcal{M})$ be the set of order filters on $\langle \mathcal{M}, \leq \rangle$ with reverse inclusion – so that $\{1\}$ is the greatest element and $\mathcal{M} = \mathbf{0}$ is the least. Union and intersection give meet and join.

We define $\nu: \mathcal{L} \rightarrow \mathcal{D}$ by

$$\nu(x) = \mathcal{M}^x.$$

This preserves order as $p \geq x \geq y$ and $p \in \mathcal{M}$ implies $p \in \mathcal{M}^y$, ie $\mathcal{M}^x \subseteq \mathcal{M}^y$.

It is one-one as $x = \bigwedge \mathcal{M}^x$.

It preserves join as $\mathcal{M}^{x \vee y} \subseteq \mathcal{M}^x \cap \mathcal{M}^y$ – by the above. And if $p \geq x$ and $p \geq y$ then $p \geq x \vee y$, so we have equality.

It preserves all meets that exist. Clearly we have $\mathcal{M}^x \cup \mathcal{M}^y \subseteq \mathcal{M}^{x \wedge y}$. And if $p \geq x \wedge y$ is meet-irreducible, then $p = (p \vee x) \wedge (p \vee y) = \bigwedge \mathcal{M}^{p \vee x} \wedge \bigwedge \mathcal{M}^{p \vee y}$. If p is in neither set then $\mathcal{M}^{p \vee x} \cup \mathcal{M}^{p \vee y} \subseteq \mathcal{M}^{>p}$ and we know that $\bigwedge \mathcal{M}^{>p} > p$ – giving a contradiction. Hence p is in one of these two sets, ie $p \geq x$ or $p \geq y$.

We can extend these ideas, a little, to elements of $\mathcal{D}$.

Definition 2.3. *Let $F \in \mathcal{D}$ and $a \in A$. Then $f_a = \bigwedge (F \cap [a, 1])$.*

Lemma 2.4. *Let $p \in \mathcal{M}$, $F \in \mathcal{D}$ and $p \geq f_a(F)$. Then $p \in F$.*

Proof. We must have $\mathcal{M}^{f_a(F)} = F \cap [a, 1]$ as $[a, 1]$ is isomorphic to $\mathcal{D}([a, 1])$ and this describes that isomorphism. $\square$

Lemma 2.5. *Let $F, G \in \mathcal{D}$ and $a \in A$. Then*

$$f_a(F \cup G) = f_a(F) \wedge f_a(G)$$

$$f_a(F \cap G) = f_a(F) \vee f_a(G).$$

Proof. For the second we have $p \geq f_a(F) \vee f_a(G)$ iff $p \geq f_a(F)$ and $p \geq f_a(G)$ iff $p \in F \cap [a, 1]$ and $p \in G \cap [a, 1]$ iff $p \in F \cap G \cap [a, 1]$ iff $p \geq f_a(F \cap G)$.

For the first we have $p \geq f_a(F) \wedge f_a(G)$ iff $p \geq f_a(F)$ or $p \geq f_a(G)$ (as $x \mapsto \mathcal{M}^x$ preserves meets) iff $p \in F \cap [a, 1]$ or $p \in G \cap [a, 1]$ iff $p \in (F \cap [a, 1]) \cup (G \cap [a, 1])$ iff $p \in (F \cup G) \cap [a, 1]$ iff $p \geq f_a(F \cup G)$. $\square$

Lastly we observe that this construction gives a universal inclusion into a distributive lattice.

Proposition 2.6. *Let $\phi: \mathcal{L} \rightarrow \mathcal{S}$ where $\mathcal{S}$ is any distributive lattice, ϕ preserves joins, one and any available meets. Then there is a unique lattice homomorphism $\hat{\phi}: \mathcal{D} \rightarrow \mathcal{S}$ such that $\phi = \hat{\phi}\nu$, ie the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{\phi} & \mathcal{S} \\ & \searrow \nu & \nearrow \iota \\ & \mathcal{D} & \end{array}$$

Proof. Define $\hat{\phi}: \mathcal{D} \rightarrow \mathcal{S}$ by

$$\hat{\phi}(F) = \bigwedge \phi[F].$$

Firstly we see that $\hat{\phi}\nu(x) = \bigwedge \phi[\mathcal{M}^x] = \phi(\bigwedge \mathcal{M}^x) = \phi(x)$ (as ϕ preserves extant meets). It is clear that $\hat{\phi}$ preserves order.

$$\hat{\phi}(F \cup G) = \bigwedge \phi[F \cup G] = \bigwedge (\phi[F] \cup \phi[G]) = \bigwedge \phi[F] \wedge \bigwedge \phi[G] = \hat{\phi}(F) \wedge \hat{\phi}(G).$$

$$\begin{aligned} \hat{\phi}(F) \vee \hat{\phi}(G) &= \bigwedge \phi[F] \vee \bigwedge \phi[G] = \bigwedge_{\substack{f \in F \\ g \in G}} \phi(f) \vee \phi(g) = \\ &= \bigwedge_{\substack{f \in F \\ g \in G}} \phi(f \vee g) = \bigwedge_{\substack{f \in F \\ g \in G}} \phi(\bigwedge \mathcal{M}^{f \vee g}) = \bigwedge_{\substack{f \in F \\ g \in G}} \phi(\mathcal{M}^{f \vee g}). \end{aligned}$$

Now $\hat{\phi}(F \cap G) = \bigwedge_{p \in F \cap G} \phi(p)$ and the two sets we're taking meets over are the same as if $p \in \mathcal{M}$ then $p \in F \cap G$ iff there are $f \in F$ and $g \in G$ with $p \geq f \vee g$.

Hence $\hat{\phi}$ is a lattice homomorphism. It might not preserve $\mathbf{0}$ as we do not know that $\bigwedge \phi[\mathcal{M}] = \mathbf{0}_{\mathcal{S}}$. $\square$

3. THE VALUATION RING

Now we turn to another construction, that of the valuation ring of a distributive lattice. We assume, as above, that $\mathcal{L}$ is a finite strong upper semilattice that is also distributive.

Definition 3.1. *For any abelian group $\mathcal{G}$, a function $f: \mathcal{L} \rightarrow \mathcal{G}$ is a valuation iff for any $x, y, z \in \mathcal{L}$*

$$f(x \vee y \vee z) + f((x \vee z) \wedge (y \vee z)) = f(x \vee z) + f(y \vee z).$$

This definition gives the usual definition if applied to a lattice – take $z = x \wedge y$.

We now consider the construction of the valuation ring of $\mathcal{L}$. This parallels the development in [4].

Now let $\mathbb{Z}^{\mathcal{L}}$ be the free abelian group over $\mathcal{L}$. This has a multiplication induced by $\langle x, y \rangle \mapsto x \vee y$ on $\mathcal{L}$ and extended by linearity.

We can define a submodule $M(\mathcal{L})$ via generators $a \vee b \vee c + (a \vee c) \wedge (b \vee c) - a \vee c - b \vee c$ for all $a, b, c \in \mathcal{L}$. This is in fact an ideal of the ring when $\mathcal{L}$ is distributive – as $t \in \mathcal{L}$ and

$a \vee b \vee c + (a \vee c) \wedge (b \vee c) - a \vee c - b \vee c \in M(\mathcal{L})$ implies

$$\begin{aligned}
 & (a \vee b \vee c + (a \vee c) \wedge (b \vee c) - a \vee c - b \vee c) \vee t \\
 &= (a \vee b \vee c \vee t) + ((a \vee c) \wedge (b \vee c)) \vee t - (a \vee c \vee t) - (b \vee c \vee t) \\
 &= (a \vee t) \vee (b \vee t) \vee (c \vee t) + ((a \vee t) \vee (c \vee t)) \wedge ((b \vee t) \vee (c \vee t)) \\
 &\quad - (b \vee t) \vee (c \vee t) - (a \vee t) \vee (c \vee t) \\
 &\in M(\mathcal{L}).
 \end{aligned}$$

On quotienting out by this subgroup we get the valuation ring $V(\mathcal{L})$.

Remark 3.1. In lattices, this reduces to the same as the usual definition for valuations.

There is a natural mapping $\iota: \mathcal{L} \rightarrow V(\mathcal{L})$ given by $x \mapsto x$. This is a valuation by the definition of $M(\mathcal{L})$. Furthermore it is a universal valuation.

Theorem 3.2. *Let $f: \mathcal{L} \rightarrow \mathcal{G}$ be any valuation. Then there is a unique group homomorphism $\phi: V(\mathcal{L}) \rightarrow \mathcal{G}$ such that the following diagram commutes:*

$$\begin{array}{ccc}
 \mathcal{L} & \xrightarrow{f} & \mathcal{G} \\
 & \searrow \iota & \nearrow \phi \\
 & V(\mathcal{L}) &
 \end{array}$$

Proof. Define $\Phi \upharpoonright \mathcal{L}$ to be f , and extend linearly to all of $\mathbb{Z}^{\mathcal{L}}$. Then Φ is zero on $M(\mathcal{L})$ – as f is a valuation – and so factors through $V(\mathcal{L})$ to give ϕ .

Uniqueness is immediate from our definition – we must have agreement with f and linearity. $\square$

Corollary 3.3. *Let $f: \mathcal{L}_1 \rightarrow \mathcal{L}_2$ be a homomorphism of suitable type (ie it preserves $\vee, \mathbf{1}$ and such meets as exist). Then there is a ring homomorphism $V(f): V(\mathcal{L}_1) \rightarrow V(\mathcal{L}_2)$ such that*

$$\begin{array}{ccc}
 \mathcal{L}_1 & \xrightarrow{f} & \mathcal{L}_2 \\
 \downarrow \iota_1 & & \downarrow \iota_2 \\
 V(\mathcal{L}_1) & \xrightarrow{V(f)} & V(\mathcal{L}_2)
 \end{array}$$

commutes.

Proof. We note that $\iota_2 f$ is a valuation on $\mathcal{L}_1$ and so the theorem gives the function $V(f)$. $\square$

This corollary shows that V is functorial. It is not hard to see that $V(f)$ is actually a ring homomorphism.

We will assume that the reader knows the basics of valuation theory on lattices in what follows. A good reference is [5].

Some properties of ι are now evident. As constant functions are valuations we see that if $g \in \mathcal{G}$ is any element then $c_g: \mathcal{L} \rightarrow \mathcal{G}$ is a valuation and so for any $l \in \mathcal{L}$ we have that the order of $\iota(l)$ is a multiple of the order of $\phi_c \iota(l) = g$. Hence $\iota(l)$ has infinite order.

We also see that if $\mathcal{L}$ is a distributive lattice then any two elements may be separated by a prime filter $\mathcal{F}$. The characteristic function of $\mathcal{F}$ into $\mathbb{Z}_2$ is a valuation and the two elements under consideration have distinct values. Hence they must be separated by ι .

Proposition 3.4. *Let $\mathcal{L}$ be a strong upper semilattice with $\mathbf{1}$ that is locally principal, ie there is a set A of elements that are all minimal in $\mathcal{L}$ such that $\mathcal{L} = \bigcup_{a \in A} [a, \mathbf{1}]$.*

Then ι is one-one on $\mathcal{L}$.

Proof. The proof proceeds by considering the distributive lattices $[a, \mathbf{1}]$ for $a \in A$ and the mappings $\iota_a: [a, \mathbf{1}] \rightarrow V([a, \mathbf{1}])$, which we already know to be one-one.

Now we have $\mathbb{Z}^{([a, \mathbf{1}])} \subseteq \mathbb{Z}^{(\mathcal{L})}$ and if t is a generator of $M([a, \mathbf{1}])$ then it is also in $M(\mathcal{L})$. Furthermore if s is a generator of $M(\mathcal{L})$ with $s \in \mathbb{Z}^{([a, \mathbf{1}])}$ then the freeness of the module implies s is a generator for $M([a, \mathbf{1}])$. Thus we have

$$M([a, \mathbf{1}]) = M(\mathcal{L}) \cap \mathbb{Z}^{([a, \mathbf{1}])}.$$

This means we have an injective homomorphism $\vartheta_a: V([a, \mathbf{1}]) \rightarrow V(\mathcal{L})$ defined by

$$\vartheta_a([x]_a) = [x]_{\mathcal{L}}$$

(the subscripts indicate where the equivalence class lives). We have the following properties of ϑ_a :

Well-defined & one-one: as $c \sim_a d$ iff $c - d \in M([a, \mathbf{1}])$ iff $c - d \in M(\mathcal{L}) \cap \mathbb{Z}^{([a, \mathbf{1}])}$ iff $c \sim_{\mathcal{L}} d$.

Preserves +: as the operation $+$ is the same in $\mathbb{Z}^{([a, \mathbf{1}])}$ as in $\mathbb{Z}^{(\mathcal{L})}$.

Preserves $\times$: as $[a, \mathbf{1}]$ is a sub-semilattice of $\mathcal{L}$ and so multiplication in $\mathbb{Z}^{([a, \mathbf{1}])}$ is the same as in $\mathbb{Z}^{(\mathcal{L})}$.

Now define a function $j_a: \mathcal{L} \rightarrow V([a, \mathbf{1}])$ by $j_a(y) = \iota_a(y \vee a)$. Then j_a is a valuation – as

$$\begin{aligned} j_a(x \vee y \vee z) + j_a((x \vee z) \wedge (y \vee z)) &= \iota_a(x \vee y \vee z \vee a) + \iota_a([(x \vee z) \wedge (y \vee z)] \vee a) \\ &= \iota_a((x \vee a) \vee (y \vee a) \vee (z \vee a)) + \iota_a((x \vee z \vee a) \wedge (y \vee z \vee a)) \\ &= \iota_a((x \vee a) \vee (y \vee a) \vee (z \vee a)) + \iota_a((x \vee a) \vee (z \vee a)) \wedge ((y \vee a) \vee (z \vee a)) \\ &= \iota_a((x \vee a) \vee (z \vee a)) + \iota_a((y \vee a) \vee (z \vee a)) \\ &= \iota_a(x \vee z \vee a) + \iota_a(y \vee z \vee a) \\ &= j_a(x \vee z) + j_a(y \vee z). \end{aligned}$$

Suppose that $\iota_{\mathcal{L}}(x) = \iota_{\mathcal{L}}(y)$.

Let $\varphi_a: V(\mathcal{L}) \rightarrow V([a, \mathbf{1}])$ be such that $j_a = \varphi_a \iota_{\mathcal{L}}$. Then $\iota_{\mathcal{L}}(x) = \iota_{\mathcal{L}}(y)$ implies $j_a(x) = j_a(y)$ and so $\iota_a(x) = \iota_a(x \vee a) = \iota_a(y \vee a)$. As ι_a is one-one we have $x = y \vee a$. Likewise $y = x \vee a$. And so $y = x \vee a = y \vee a \vee a = y \vee a \geq a$. (and $x \geq a$.)

But now $\vartheta_a \iota_a(x) = \iota_{\mathcal{L}}(x) = \iota_{\mathcal{L}}(y) = \vartheta_a \iota_a(y)$ implies $x = y$ – as both ϑ_a and ι_a are one-one. $\square$

In this result we have a lattice homomorphism $\text{incl}: [a, \mathbf{1}] \rightarrow \mathcal{L}$ and it's clear that $\vartheta_a = V(\text{incl})$. Because $\mathcal{L}$ is distributive, the mapping $J_a: x \mapsto x \vee a$ is a lattice homomorphism from $\mathcal{L} \rightarrow [a, \mathbf{1}]$ and $\varphi_a = V(J_a)$. Also $\text{incl} \circ J_a = \text{id}_a$ and so $\varphi_a \vartheta_a = V(\text{id}_a) = \text{id}$ and so we actually have a retract. This gives a kind of localization on $V(\mathcal{L})$ that is extremely useful.

3.1. The envelope. Now that we have the valuation ring we can define a new partial operation that produces our envelope.

We begin by defining

$$x \wedge y = x + y - x \vee y$$

and extending for any set $X \subseteq \mathcal{L}$ by letting $X_{(m)} = \{x \mid x \text{ is minimal in } X\}$.

$$\bigwedge X = \sum_{\substack{B \subseteq X_{(m)} \\ B \neq X_{(m)}}} (-1)^{|B|+1} \bigvee B.$$

The envelope is the range of this operation. We want to show that this indeed produces a distributive lattice.

Rather than go through the tedious calculations involved we consider the relationship between the valuation ring of $\mathcal{L}$ and the valuation ring of the distributive envelope constructed above, ie $V(\mathcal{D})$.

4. THE TWO ENVELOPES ARE EQUIVALENT

We want to consider how to extend valuations on $\mathcal{L}$ to valuations on $\mathcal{D}$. Of course we need only extend the universal valuation ι . The most natural method is to equate a filter $F \in \mathcal{D}$ with its “virtual meet” $\bigwedge F = \bigwedge_{a \in A} f_a(F)$ and use inclusion-exclusion to define

$$i(F) = \sum_{\substack{B \subseteq A \\ |B| \geq 1}} (-1)^{|B|+1} \bigvee_{b \in B} f_b(F).$$

However it might be simpler to just use all of F and define it as

$$j(F) = \sum_{\substack{H \subseteq F \\ |H| \geq 1}} (-1)^{|H|+1} \bigvee H.$$

The calculations involved in checking this are not easy and so we argue indirectly that this method must work.

We know that valuations on a distributive lattice are completely determined by their values on the set of meet-irreducibles and $\mathbf{1}$ (because these form a basis). Furthermore we know that (for distributive lattices) that we can define a valuation from any set of values on the meet-irreducibles and $\mathbf{1}$ (same reason). Putting these together means that we can start with a valuation on $\mathcal{L}$, lift to one on $\mathcal{D}$ by using its values on $\mathcal{M}(\mathcal{L}) = \mathcal{M}(\mathcal{D})$, and restrict to $\mathcal{L}$.

Now the inclusion/exclusion lemma holds true in both $V(\mathcal{L})$ and in $V(\mathcal{D})$.

Lemma 4.1. *Let $x_1, \dots, x_n \in \mathcal{L}$ be such that $\bigwedge_i x_i$ exists. Then*

$$\bigwedge_i x_i = \sum_{\substack{J \subseteq \{1, \dots, n\} \\ |J| > 0}} (-1)^{|J|+1} \bigvee_{j \in J} x_j.$$

Proof. Let $a \leq \bigwedge_i x_i$ be minimal. Then it holds in $V([a, \mathbf{1}])$ which is a subalgebra of $V(\mathcal{L})$ and so it holds in $V(\mathcal{L})$. $\square$

By this inclusion/exclusion formula in $V(\mathcal{L})$ the restriction must be same as the valuation we started with. This works both ways: as if f is a valuation on $\mathcal{D}$ then it restricts to one on $\mathcal{L}$. Thus the valuation rings must be the same – ie this mapping makes both $V(\mathcal{D})$ and $V(\mathcal{L})$ universal valuations for $\mathcal{D}$ and so they are isomorphic.

Thus either of the definitions above must work, and be the same. In particular, if $X \subseteq \mathcal{L}$ and F_X is the filter generated by X then $\bigwedge X = F_X$ in $\mathcal{D}$ and we see that X and F_X have the same set of minimal elements $X_{(m)}$. Therefore $\bigwedge X_{(m)} = F_X$ in $\mathcal{D}$ also. Then

$$\bigwedge X = \sum_{\substack{B \subset X_{(m)} \\ B \neq X_{(m)}}} (-1)^{|B|+1} \bigvee B = \bigwedge F_X = F_X.$$

Thus the envelope we defined above is exactly $\mathcal{D}$. This gives us our alternative representation for $\mathcal{D}$.

REFERENCES

- [1] J. C. Abbott, *Algebras of implication and semi-lattices*, Séminaire Dubreil. Algèbre et théorie des nombres **20**, 1–8.
- [2] J. C. Abbott, *Sets, Lattices, and Boolean Algebras*, Allyn and Bacon, Boston, MA, 1969.
- [3] I. Chajda, R. Halaš, and J. Kühr, *Implication in MV-algebras*, Algebra Universalis **52** (2004), 377–382.
- [4] M. Barnabei, A. Brini, and G.-C. Rota, *The theory of Möbius functions*, Uspekhi Mat. Nauk. **41** (1986), 113–157.
- [5] L. Geissinger, *Valuations on Distributive Lattices I*, Arch.Math. **24** (1973), 230–239.

SCHOOL OF MATHEMATICS, STATISTICS AND OPERATIONS RESEARCH, VICTORIA UNIVERSITY OF WELLINGTON, WELLINGTON, NEW ZEALAND,

E-mail address: Colin.Bailey@vuw.ac.nz

PACIFIC NORTHWEST NATIONAL LABORATORIES, RICHLAND, WA, U.S.A.

E-mail address: Joseph.Oliveira@pnl.gov